



Contribution ID : 61

Type : Oral

Shape evolution and Collectivity beyond 78Ni : Lifetime measurements of low-lying states in neutron-rich Zn

Tuesday, 18 July 2023 16:25 (15)

Nuclear shape is a sensitive probe of understanding the many-body quantum system and nucleon-nucleon interaction. Shape coexistence was reported in doubly magic 78Ni [1]. Related features such as triaxiality [2] and onset of deformation beyond $N = 50$ [3,4] were also reported in this mass region. The study of these phenomena plays a crucial role in understanding the limit of nuclear stability as well as the predicted fifth island of inversion [5]. One of the observables experimentally to study nuclear shape is the lifetime of excitation state, which has a direct link with the electric quadrupole moment Q . In a recent gamma spectroscopy study of $82,84\text{Zn}$ [4], the magicity was confined to $N = 50$ in 80Zn only, while an onset of deformation for low-lying states was identified with the help of $E(2^+)/E(41^+)$ and $E(21^+)/E(41^+)$ ratios towards heavier Zn isotopes. However, the lifetimes of these states are still unknown. Therefore, lifetime measurement of low-lying states was performed in neutron-rich Zn isotopes to further investigate the shape evolution and development of collectivity beyond $N = 50$.

Neutron-rich Zn isotopes were investigated at RIKEN Nishina Center during the HiCARI 2020 campaign. 345 MeV/u ^{238}U impinged on ^9Be primary target with an average intensity 60 pA. Production fragments were then separated and identified by BigRIPS spectrometer. A secondary 6 mm thick ^9Be target was placed at F8 to induce knockout reactions. After the target, ion of interests were identified on an event-by-event base by using $B\rho-\Delta E$ -TOF technique with the ZeroDegree spectrometer. The secondary target was surrounded by HiCARI, consisting of 6 Miniball triple clusters, 4 Clovers and 2 Gretina-type tracking clusters, which was used for Doppler correction of gamma rays from in-flight ions and lifetime measurement.

Some low-lying states in neutron-rich $76-82\text{Zn}$ were established based on the recent experiment. The lifetime of each state was determined by gamma-ray lineshape analysis [6]. The shape evolution in neutron-rich Zn isotopes will be discussed by comparing the experimental results with shell model and mean-field calculations.

R. Taniuchi et al., Nature 569, 53 (2019).

A. M. Forney et al., PRL 120, 212501 (2018).

C. Delafosse et al., PRL 121, 192502 (2018).

C. Shand et al., PLB 773, 492 (2017).

F. Nowacki et al., PRL 117, 272501 (2016).

P. Doornenbal et al., NIMA 613, 218 (2010).

Primary author(s) : Dr. CHEN, Zhiqiang (GSI); Prof. PODOLYÁK, Zsolt (University of Surrey); Prof. FLAV-IGNY, Freddy (LPC Caen); Dr. GÓRSKA-OTT, Magdalena (GSI); COLLABORATION, RIBF196

Presenter(s) : Dr. CHEN, Zhiqiang (GSI)

Session Classification : Session 7A

Track Classification : Experimental Nuclear Structure