



Contribution ID : 66

Type : Oral

Cross sections for $^{54}\text{Fe}(n,n')$ ^{54}Fe and $^{54}\text{Fe}(n,p')$ ^{54}Mn deduced from the detection of de-excitation γ rays

Tuesday, 18 July 2023 17:10 (15)

Iron is an important component of many structural materials; examples include energy production complexes, laboratories, devices, and shipping containers that often cross borders. The properties of iron alloys in the structural materials—strength, ductility, and stability—depend on defects that develop and grow from neutron scattering and (n, p) and (n, γ) reaction rates. ^{54}Fe is only 5.5% abundant, but neutron scattering cross sections and reaction rates for this nucleus can affect fast reactor systems and energy transport and deposition. Neutron scattering cross sections obtained by the detection of the scattered neutrons offers the clearest path to the desired cross sections in the fast neutron region¹, but such measurements are typically limited to scattering from only the lowest few excited levels because of the large energy spreads (10s to 100s of keV) inherent in neutron experiments. The detection of de-excitation γ rays (< 2.5 keV resolution) following inelastic scattering or proton production on ^{54}Fe offers a rare opportunity to investigate (n, p) and (n, n') cross sections to higher-lying levels in a consistent way by the examination of γ -ray production rates.

Measurements have been performed on an enriched ^{54}Fe sample (97.6%) using the neutron production and γ -ray detection facilities at the University of Kentucky Accelerator Laboratory. γ -ray excitation functions were measured for incident neutron energies from 1.5 to 4.7 MeV. Angular distributions and Doppler shifts were measured at $E_n = 4.5$ MeV. γ -ray production cross sections were deduced by considering all γ rays feeding or resulting from the decay of each level. Analogous measurements were made on natural Ti, V, Al and Fe samples for absolute normalization purposes. The results of our measurements will be presented in comparison to evaluated data (ENDF, JENDL, JEFF libraries) and with TALYS calculations.

This work was supported by the U.S. Department of Energy NNSA-SSAP award NA-0002931 and Nuclear Energy Universities Program award NU-12-KY-UK-0201-05, the U.S. National Science Foundation under grant PHY-1305801, and the U.S. National Isotope Development Program.

[1] J. R. Vanhoy et al., Nucl. Phys. A 972, 107 (2018).

Primary author(s) : HICKS, Sally (University of Kentucky; University of Dallas); PECHA, R. L. (University of Dallas); HOWARD, Thaddeus (University of Dallas); FRENCH, A. J. (University of Dallas); SANTONIL, Z. C. (University of Dallas); VANHOY, J. R. (United States Naval Academy); RAMIREZ, A. P. D. (University of Kentucky); PETERS, E. E. (University of Kentucky); LIU, S. H. (University of Kentucky); PRADOS-ESTÉVEZ, F. M. (University of Kentucky); ROSS, T. J. (University of Kentucky); CRIDER, B. P. (Mississippi State University); YATES, S. W. (University of Kentucky)

Presenter(s) : HICKS, Sally (University of Kentucky; University of Dallas)

Session Classification : Session 7B

Track Classification : Nuclear Reactions